

Scientific Note

Genetic Structure of Three Populations of
Oxya chinensis in Shanxi, ChinaHAN Yan¹, DUAN Yi-hao², MA En-bo¹, QIAO Hai-xuan¹

(1. Department of Life Science; 2. Department of Environmental Science, Shanxi University, Taiyuan 030006, China)

Abstract: Starch gel electrophoresis was used to analyze genetic structure of three populations of *Oxya chinensis* from Shanxi Province, China. Four allozyme polymorphic loci (MDH-1, MDH-2, LDH, ME) were found. The data suggest that two populations of *O. chinensis* about 41 kilometers apart are genetically similar, but differentiated from the third population with a geographic distance of 357 kilometers. However, the heterozygosity levels of those two genetically similar populations were much different. The Nei's genetic distance between two geographically close populations is 0.068, which is much smaller than those of geographically distant populations (0.23 in average). The observed overlap of geographical distance and genetic distance implies that for *O. chinensis* population divergence may correlate with geographic isolation.

Key words: *Oxya chinensis*; Allozyme; Population genetics; Shanxi

Received date: 2001-06-18; Accepted date: 2001-08-31

Foundation item: The project was supported by the National Natural Science Foundation of China (30070112)

收稿日期: 2001-06-18; 接受日期: 2001-08-31

(上接第 75 页)

Evol., 3: 254-259.

Ocarido M, Siefert L, Baranga J. 1999. Seasonal changes of impala (*Aepyceros melampus* Lichtenstein, 1812) faecal helminth egg counts through a one-year period [J]. *African Journal of Ecology*, 37(3): 327-333.

Pei Y X. 2000. The effect of diet quality on the passage rate, digestive performance, and gastrointestinal tract morphology in two rodent species from inner Mongolia grassland [D]. Master's degree dissertation of Institute of Zoology, the Chinese Academy of Sciences. [裴艳新. 2000. 食物质量对内蒙古草原两种啮齿动物的食物通过速率、消化性能及消化道形态的影响. 中国科学院动物研究所硕士毕业论文.]

Phillips M K, Schreck J. 1991. Parasitism in captive and reintroduced red alves [J]. *J. Wildl. Disease*, 27: 498-501.

Putman R J. 1984. Facts from faeces [J]. *Mammal Rev.*, 14: 79-97.

Queller D C, Strassmann J E, Hughes C R. 1993. Microsatellites and kinship [J]. *Trends Ecol. Evol.*, 8: 285-288.

Ramirez R G. 1999. Feed resources and feeding techniques of small ruminants under extensive management conditions [J]. *Small Ruminant Research*, 34(3): 215-230.

Reed J Z, Tollit D J, Thompson P M, et al. 1997. Molecular scatology, the use of molecular genetic analyses to assign species, sex and individual identity to seal faeces [J]. *Molecular Ecology*, 6: 225-234.

Robertson B C, Minot E O, Lambert D M. 1999. Molecular sexing of individual kakapo, *Strigops habroptilus* Aves, from feces [J]. *Mol. Ecol.*, 8: 1347-1350.

Rosen D A S, Trites A W. 2000. Digestive efficiency and dry-matter digestibility in Steller sea lions fed herring, pollock, squid, and salmon [J]. *Canadian Journal of Zoology*, 78(2): 234-239.

Schoenly K, Cohen J E. 1991. Temporal variation in food web structure: 16 empirical cases [J]. *Ecological Monographs*, 61(3): 267-298.

Smith A D. 1964. Defaecation rates of mule deer [J]. *Journal of Wildlife Management*, 28: 435-444.

Soderquist T R. 1995. Spatial organization of the arboreal carnivorous marsupial *Phascogale tapoatafa* [J]. *Journal of Zoology (London)*, 237(3): 385-398.

Steven J. 1992. Fighting the rules of the game [J]. *Nature*, 355: 489.

Strauch D. 1992. Environmental pollution by animal production and possible measures for their reduction: Hygienic aspects [J]. *Zuechtungskunde*, 64(3-4): 245-253.

Sugiyama Y, Kawamoto S, Takenaka O, et al. 1993. Paternity discrimination and inter-group relationships of chimpanzees at Bossou [J]. *Primates*, 34(4): 545-552.

Sutherland W J. 1996. Ecological Census Technique: A Handbook [M]. London: Cambridge University Press.

Wei F W, Rao G, Lu M, et al. 2001. Molecular scatology and its application: Reliability, limitation and prospect [J]. *Acta Theriologica*, 21(2): 143-152. [魏辅文, 饶刚, 李明, 等. 2001. 分子粪便学及其应用: 可靠性、局限性和展望. 兽类学报, 21(2): 143-152.]

White G C, Carrott R A. 1990. Analysis of Wildlife Radio-tracking Data [M]. San Diego: Academic Press.

Wu C H, Su B, Zhang Y P. 1997. Non-invasive DNA determination in several endangered species [A]. In: Jiang Z G, Ma K P, Han X G. Conservation Biology [M]. Hangzhou: Zhejiang Publishing House of Scientific and Technology. 194-197. [吴春花, 宿兵, 张亚平. 1997. 濒危动物的非损伤性 DNA 检测. 见: 蒋志刚, 马克平, 韩兴国. 保护生物学. 杭州: 浙江科学技术出版社. 194-197.]

中华稻蝗三种群遗传结构分析

韩 焱¹, 段毅豪², 马恩波¹, 乔海暄¹

(1. 山西大学 生命科学系; 2. 山西大学 环境科学系, 山西 太原 030006)

摘要: 应用淀粉凝胶电泳对山西省太原黄陵村、临猗县赵村及永济县伍姓湖 3 个中华稻蝗 (*Oxya chinensis*) 种群的 4 个等位酶位点 (MDH-1、MDH-2、LDH、ME) 进行了分析。结果表明 3 个种群的中华稻蝗在遗传结构上具有差异: 黄陵种群的 MDH-1、MDH-2 以及伍姓湖种群的 MDH-1 均为单态位点; 赵村种群的杂合性最低 ($H = 0.112$)、伍姓湖种群最高 ($H = 0.229$); 赵村种群与伍姓湖种群间的 Nei's 遗传距离为 0.068, 可视为 1 个大的种群, 而黄陵种群与赵村种群和伍姓湖种群的 Nei's 遗传距离分别为 0.247 和 0.218。考虑到赵村和伍姓湖之间的地理距离 (41 km) 比太原黄陵 (357 km) 要近得多, 结果提示 3 个种群地理距离与遗传结构差异之间存在相关关系。

关键词: 中华稻蝗; 等位酶; 种群遗传; 山西省

中图分类号: Q969.265.2; Q953 **文献标识码:** A **文章编号:** 0254-5853(2002)01-0076-05

Oxya chinensis (Orthoptera, Acrididea, Catantopidae, *Oxya* Serville) represents an important agricultural pest to rice and other crops. This grasshopper species widely and abundantly distributes in most part of China and the adjacent regions. Taxonomically *O. chinensis* was previously described as a species complex with several species involved (Ma, 1991). However, due to the subtlety of morphological characteristics used for species definition, some researchers suggested that as many as ten described species could be grouped as one species of *Oxya chinensis* (Hollis, 1971, 1975). In the past ten years the comparative studies on chromosome have showed that the C-band types of different *O. chinensis* populations across China (from Heilongjiang to Fujian Province) demonstrated very high similarity (Ma *et al.*, 1994), indicating the relatively low differentiation among *O. chinensis* populations. More recent research on the cytochrome-*b* gene in mitochondrial DNA also supports the view that the *O. chinensis* populations are less divergent than those of other grasshopper species (Wang, 2001).

However, several studies have noted quantitative morphological difference among *O. chinensis* populations. For example, there exists an increasing cline of body size of *O. chinensis* from the north to south in China. The body length of the samples from Nenjiang, Heilongjiang Province (22.45 ± 2.05 mm) is much smaller than those from Fuzhou, Fujian Province (33.65 ± 2.45 mm) (Guo *et al.*, 2001). Another research examined other morphological characters between two populations from Chang'an and Hanzhong, Shaanxi Province. The data suggest that the quantitative differences are significant enough to

define *O. chinensis* from those two sites as relatively independent different demes (Wang *et al.*, 1997). Because the body size and other quantity characters of *O. chinensis* are genetically determined and stable among the populations, it becomes necessary to examine the genetic differentiation of *O. chinensis* at molecular level in order to explore the possible evolutionary history of this species.

Allozyme analysis has been used as a valuable method to examine population genetic structure and compare differentiations. It has been successfully employed to study a wide variety of taxa. However, its application to grasshopper is scarce. Zheng *et al.* (1986) compared esterase zymograms in 8 locust species from China. Later the study was expanded by using two more isozymes of malate dehydrogenase and lactate dehydrogenase on 67 species with a few individuals per species (Huang, 1990). The present study used allozyme analysis to examine the genetic structure of three populations of *O. chinensis* from Shanxi Province in Midwest China.

The hypothesis to be tested is that the geographically isolated populations of *O. chinensis* may demonstrate genetic differentiation at allozyme level and the divergence levels may correlate to geographic distance.

1 Methods and Materials

1.1 Sample collection

Three populations of *O. chinensis* were all collected from Shanxi Province, China. They are from Huangling in suburb of Taiyuan ($E 112^{\circ}36'$, $N 37^{\circ}46'$), Zhaocun Village, Linyi County ($E 110^{\circ}24'$, $N 35^{\circ}15'$), and Wuxinghu, Yongji County ($E 110^{\circ}28'$, $N 34^{\circ}53'$), in July and August, 2000.

Huangling, Zhaoacun and Wuxinghu collection areas are a grassplot, a beach with some ruderal of the grass family, and a thick reed at Wuxinghu, respectively.

The insect samples were brought to laboratory alive and then frozen at -20°C until electrophoresis. No observable mortality occurred during the transportation.

1.2 Methods

Starch gel was prepared using a mixture of 12.5 g soluble starch, 6.25 g potato starch for electrophoresis (Sigma) and 6.25 g refined potato starch (prepared in our laboratory) per 200 mL of buffer. The final starch gel concentration is 12.5% (w/v). Tris-citric acid (0.1 mol/L, pH8.0) was used as electrode buffer. The ratio of electrode buffer to gel buffer is 9:1. Gel was prepared in a mold of 135 mm \times 135 mm \times 7 mm. The cooled gel was then stored at 4°C for 2 hours before used.

The insect tissue was from femur muscle. The tissue was removed and then homogenated in 20 μL double distilled water on an ice pan. Filter paper was cut into wicks for loading sample. Electrophoresis lasted for 4 hours at 4°C , at a voltage of 140 V. The initial current was around 25 mA. Allozyme staining was conducted following the standard method (Murphy *et al.*, 1996) with modifications. The band with the fastest migration is designated "A", the second "B", and so on. Three allozymes, which were lactate dehydrogenase (LDH), malate dehydrogenase (MDH) and malic enzyme (ME), were used to examine. The LDH substrate was prepared using lactic acid (Wuhan Dongshan Lactic Acid Factory) and adjusting pH with NaOH to a final concentration of 1 mol/L. Several days later, the pH of this sodium lactate solution dropped to as low as 6.5. It has to be readjusted back to 8 before staining.

1.3 Statistics

BIOSYS-II was used to calculate allele frequency, percent polymorphic loci, heterozygosity (H), goodness-of-fit to Hardy-Weinberg's (H-W) equilibrium, Nei's genetic identity (I) and Roger's genetic distances (D) (Nei, 1987). The genetic distance was clustered with unweighted pair-group method using arithmetic averages (UPGMA).

2 Results

2.1 Locus

Using the methods in this study, all three allozymes were successfully stained with sufficient separation and clear bands that could be scored with minimum efforts. MDH has shown two loci, one (designed as MDH-1) migrated from cathode to anode, and the other (MDH-2) moved from anode to cathode. MDH-1 has three subbands. LDH and ME each have one locus, moving from cathode to anode.

2.2 Allele

Table 1 presents the allele frequency of polymorphic loci for each population.

2.3 Heterozygosity

In three populations, heterozygosity of Zhaoacun is the lowest ($H = 0.112$) and that of Wuxinghu is the highest ($H = 0.229$) (Table 2).

2.4 H-W equilibrium

The deviation from H-W equilibrium was observed in the following loci: ME of Huangling population; all loci of Zhaoacun population; and MDH-2 and ME of Wuxinghu population. (Chi-square test, $P < 0.01$) (Table 3).

2.5 Nei's genetic identity and Roger's genetic distance

Nei's genetic identity and Roger's genetic distance (Table 4) showed that Zhaoacun population and Wuxinghu population were genetically identical, so they can be regarded as two parts of a whole population. However, the genetic identities between these two populations and Huangling population are greater ($D = 0.83 - 0.85$), a typical value for populations in a single species. Cluster based on Roger's genetic distance is presented in Fig. 1, and the increase in genetic distance along with geographical distance is shown in Fig. 2.

3 Discussion

Several alleles of all the three populations of *O. chinensis* have deviated from H-W equilibrium. In most cases, the deviation is because of heterozygote deficiency. Due to species' relatively low flight capability and the water requirement for habitats, without excluding the possible inbreeding and null alleles, population subdivision and selection may be attributed to such deviation (Duan *et al.*, 1997, 2000). For example, the Nei's genetic identity between Zhaoacun and Wuxinghu populations is as high as 0.988, suggesting that they are from the same population. However, population subdivision is indicated by their dif-

Table 1 Allele frequency of polymorphic loci among three populations of *Oxya chinensis* in Shanxi

Locus	Population		
	Huangling	Zhaocun	Wuxinghu
MDH-1			
N	53	25	24
A	0.019	0.160	0.000
B	0.953	0.820	0.958
C	0.028	0.020	0.042
MDH-2			
N	53	25	24
A	0.000	0.640	0.583
B	0.991	0.360	0.396
C	0.009	0.000	0.021
LDH			
N	50	26	26
A	0.010	0.000	0.000
B	0.750	0.808	0.808
C	0.240	0.192	0.192
ME			
N	37	28	27
A	0.216	0.214	0.278
B	0.608	0.446	0.370
C	0.176	0.339	0.352

ferences in heterozygosity and percentage of polymorphic loci (Table 2). Moreover, all the four loci in Zhaocun are deviated from H-W equilibrium, but such deviation occurs at two loci in Wuxinghu population. Therefore, it is likely that the Zhaocun's samples of *O. chinensis* is a subdivision of Wuxinghu population. Besides, the parthenogenesis has been observed in *Oxya formosana*, a sibling species of *O. chinensis* (Hong & Ando, 1998). Parthenogenesis can make the population deviate from H-W equilibrium. It is not clear whether there exists parthenogenesis in *O. chinensis*.

Chromosome analysis of *O. chinensis* indicates C-

band type has an extremely high similarity among *O. chinensis* populations (Ma *et al.*, 1994). The allozyme data, however, suggests quantitative differentiation. The different perspectives of the two methods may have contributed to this observed variance. In karyotic analysis C-band indicates heterochromatin section on chromosome constituting highly repeated DNA sequence that shows the least biological activity, and therefore is conservative (Ma *et al.*, 2000). Allozyme genes, however, represent the most active genes in organisms and are subject to founder effect and natural selection. It implies that for *O. chinensis* the differentiation showed by allozyme level may not be revealed by chromosome analysis.

The three populations of *O. chinensis* sampled in this study were from Shanxi Province with a geographic distance from 41.4 to 372 km. Noticeably, a larger genetic distance between two populations was observed as the geographic distance increased (Fig 2). It suggests a positive correlation between those two important measurements in population genetics meaning that distance-by-isolation plays an important role in population's divergence. Even so more studies are needed to reconfirm this observation with expanded sampling areas and increased number of allozymes, our results offer some implications for the relationship between geographic barrier and population genetic variation in *O. chinensis*.

The life cycle of *O. chinensis* depends on much water and the species' dispersal capacity is relatively limited in terms of flight ability when compared with some other insects of Orthoptera like *Locusta migratoria*, therefore it is reasoned that a moderate geographical

Table 2 Genetic variability of three populations of *Oxya chinensis* in Shanxi

Population	Mean sample size per locus	Mean No. of alleles per locus	Percentage of polymorphic loci *	Mean heterozygosity	
				Direct-count	H-W expected **
Huangling	48.3 ± 3.8	2.8 ± 0.3	50.0	0.181 ± 0.075	0.264 ± 0.126
Zhaocun	26.0 ± 0.7	2.5 ± 0.3	100.0	0.112 ± 0.052	0.437 ± 0.081
Wuxinghu	25.3 ± 0.8	2.5 ± 0.3	75.0	0.229 ± 0.103	0.396 ± 0.108

* A locus is considered polymorphic if the frequency of the most common allele does not exceed 0.95; ** Unbiased estimate (Nei, 1987).

Table 3 Chi-square test for deviation from H-W equilibrium in three populations of *Oxya chinensis* in Shanxi

	MDH-1	MDH-2	LDH	ME
Huangling	0.103	—	1.142	19.318 *
Zhaocun	14.115 *	26.312 *	16.312 *	24.744 *
Wuxinghu	0.022	25.107 *	0.023	12.026 *

* $P < 0.01$

Table 4 The genetic similarity and distance in three populations of *Oxya chinensis* in Shanxi

	Huangling	Zhaocun	Wuxinghu
Huangling	—	0.247	0.218
Zhaocun	0.835	—	0.068
Wuxinghu	0.857	0.988	—

Below diagonal is Nei's genetic identity and above diagonal is Rogers' genetic distance.




Fig. 1 Cluster analysis of based on Roger's genetic distance in three populations of *Oxya chinensis* in Shanxi (Cophenetic coefficient = 0.997)

barrier may significantly impede the gene flow among *O. chinensis* populations, and high genetic differentiation between the populations of *O. chinensis* may result from distance-by-isolation for a long time, which quantitatively supports our hypothesis that the geographically isolated populations of *O. chinensis* demonstrate genetic




Fig. 2 Relationship between genetic distance and geographic distance in three populations of *Oxya chinensis* in Shanxi

1: the distance between Zhaoxun and Wuxinghu; 2: distance between Huangling and Zhaoxun; 3: distance between Huangling and Wuxinghu.

differentiation at allozyme level and the divergence levels may correlate to geographic distance.

Acknowledgements: We thank LI Chun-xuan, HUANG Deng-yu and SHEN Yong-ming for their assistance of collecting *Oxya* samples.

References:

- Duan Y, Guttman S I, Oris J T. 1997. Genetic differentiation among laboratory populations of *Hyalella azteca*: implications for toxicology[J]. *Environmental Toxicology and Chemistry*, **16**: 691 - 695.
- Duan Y, Guttman S I, Oris J T, et al. 2000. Genetic structure and relationships among populations of *Hyalella azteca* and *H. montezuma* (Crustacea: Amphipoda)[J]. *Journal of North American Benthological Society*, **19**: 308 - 320.
- Guo Y P, Duan Y H, Bai G R, et al. Lack of chromosomal polymorphism among *Oxya chinensis* population of different body size in China[J]. *Acta Zoologica Sinica*, (in press)
- Hollis D. 1971. A preliminary of the genus *Oxya* Audinet-Serville (Orthoptera: Acridoidea)[J]. *Bull. Br. Mus. (Nat. Hist.) Ent.*, **26** (7): 269 - 343.
- Hollis D. 1975. A revision of the subfamily Oxyinae (Orthoptera: Acridoidea)[J]. *Bull. Br. Mus. (Ent.)*, **31**: 191 - 234.
- Hong Z D, Ando Y. 1998. Parthenogenesis in three species of genus *Oxya* [J]. *Jap. J. Applied Ent. & Zool.*, **42**(2): 65 - 69.
- Huang Y. 1990. Chemotaxonomic study of Acridoidea insects in China (Orthoptera)[D]. Ph. D. dissertation of Shaanxi Normal University. 103 - 110. (in Chinese).
- Ma E B. 1991. Cytotaxonomic study of catantopidae in China (Orthoptera: Acridoidea)[D]. Ph. D. dissertation of Shaanxi Normal University. 233 - 234. (in Chinese).
- Ma E B, Guo Y P, Zheng Z M. 1994. Cytotaxonomic study of *Oxya* species in China (Orthoptera: Acridoidea)[J]. *J. Entomologica Sinica*, **1** (2): 101 - 109. (in Chinese).
- Ma E B, Guo Y P, Ren Z M, et al. 2000. Chromosomal C-banding karyotype in three species of *Oxya* and its cytotaxonomic relationship [A]. In: Li D M, Zhang Z L, Huang D W, et al. Proceedings of Annual Conference of Chinese Entomological Society [C]. Beijing: China Science and Technology Press. 10: 59 - 62. (in Chinese).
- Murphy R W, Sites J W, Butth D J, et al. 1996. Proteins I: Isozyme electrophoresis [A]. In: Hillis D M, Moritz C, Mable B K. Molecular Systematics (2nd edition) [M]. Sunderland, MA, USA: Sinauer Associates, Inc. Publisher. 114.
- Nei M. 1987. Molecular Evolutionary Genetics [M]. New York: Columbia University Press.
- Wang B C. 2001. Mitochondrial DNA sequences analysis and phylogenetic studies on some species of genus *Oxya* in China [D]. Master's degree dissertation of Shanxi University. 29. (in Chinese).
- Wang H C, Wang Q C, Liu Z B, et al. 1997. Comparative study on two populations of *Oxya chinensis* (Thurberg) in Shaanxi province [J]. *Acta Entomologica Sinica*, **40**(4): 374 - 377. (in Chinese).
- Zheng Z M, Xiong Z Y, Lian Z M. 1986. EST isozyme research of eight species grasshopper [J]. *Journal of Shaanxi Normal University*, (11): 52 - 59. (in Chinese).